

# CHILDHOOD IMMUNISATION SCHEDULE

**ROUTINE VACCINES**  
in the Singapore's  
Health Promotion Board  
immunisation schedule.†

AGE	WEEK	MONTH	YEAR
BCG	0 wk		
Hep B	0 wk	1 mth	6 mth
DTP/Tdap		3 mth 4 mth 5 mth	18 mth 10-11 yr
OPV/IPV		3 mth 4 mth 5 mth	18 mth 6-7 yr 10-11 yr
MMR			12 mth 15-18 mth
Pneumococcal		3 mth 5 mth	12 mth
DTPa-IPV/Hib (5-in-1)	Hep B 0 wk	Hep B 1 mth 3 mth 4 mth 5 mth	Hep B 6 mth 5-in-1 18 mth
DTPa-IPV/Hib/HepB (6-in-1)	Hep B 0 wk	1.5, 3, 5 mth or 2, 4, 6 mth 5-in-1 18 mth	
DTPa-IPV/Hib/HepB (modified)	Hep B 0 wk	Hep B 1 mth 5-in-1 3 mth 5-in-1 4 mth 6-in-1 5 mth	5-in-1 18 mth
MMRV			12 mth 15-18 mth
Rotavirus 2 or 3 doses depending on brand used. See notes.		6 wk 3 mth	
Hib		3 mth 4 mth 5 mth	18 mth
Varicella			2 doses given 6-10 weeks apart
Hep A			After 1 yr: 2 doses 6 mths apart
Influenza			From 6 mths: 2 doses 1mth apart for first time vaccination (< 9 yrs old); thereafter 1 dose every 6 mths to a year
HPV 2 or 3 doses depending on brand used. See notes.			9-26 yrs 3 doses: 0, 1/2, 6 mth

**COMBINATION VACCINES**

**RECOMMENDED VACCINES**

Please refer to full prescribing information before prescribing. Reference: 1) Immunisation Schedule, 1st Dec 2011. Health Promotion Board, Singapore.

**Primary dose**  
**Booster dose**

**BCG** Ideally given at birth, if given beyond 12 mths, do PPD. Give BCG if negative.  
**Hep B** The birth dose may be used as the 1st dose in a 3 dose primary series with 4 wk intervals. Neonates of carrier mothers: 4 doses at 0, 1, 2 & 12 mths or 3 doses at 0, 1 & 6 mths.

**DTP/Tdap** 2nd Booster for DTP is given as reduced antigen Tdap.  
**Rotavirus** Doses from 6 wks to 6 mths, 2 or 3 doses, and all doses must be completed by 6 mths or 32 wks of age depending on brand used. 1st dose preferably given to infants between 6-14 wks of age. Interval between doses should not be less than 4 wks. OPV and RV doses must be given 2 wks apart.

**Pneumococcal** Children < 12 mths: 2 doses + 1 booster, 12 - 23 mths: 2 doses, 24 - 59 mths: 1 dose  
**HPV** Cervarix (10-25 yrs): 3 doses at 0, 1, 6 mths. Gardasil (9-26 yrs): 3 doses at 0, 2, 6 mths.